

Oj boli, boli, czyli ból w okresie okooperacyjnym chirurgii soczewki

Oh, It Hurts – Perioperative Pain in Cataract Surgery

Ewa Langwińska^{1,2}, Karolina Pielak², Agata J. Ordon^{2,3}, Katarzyna Malinowska²

¹ Katedra i Klinika Okulistyczna Wydziału Lekarskiego Warszawskiego Uniwersytetu Medycznego

Kierownik: prof. dr hab. n. med. Jacek P. Szaflik

² Samodzielny Publiczny Kliniczny Szpital Okulistyczny w Warszawie

³ Centrum Mikrochirurgii Oka LASER w Warszawie

Summary:

Pain and anesthesia are important parts of the perioperative management of patients undergoing cataract surgery. They affect the quality of life and the readiness to undergo another procedure.

The level of anxiety, pain, and its intensity occur with different frequencies.

The group of patients who underwent uncomplicated phacoemulsification cataract surgery under local anesthesia and intravenous sedation was asked to complete an anonymous questionnaire, taking into account several factors such as age, gender, level of education, level of anxiety before surgery, nature of pain, previous procedures, general diseases and other.

The study aims to compare the frequency and intensity of perioperative pain during first and second eye surgery and to list predisposing factors to pain during cataract surgery.

Key words:

cataract, pain, pain perception, anxiety, phacoemulsification, cataract surgery.

Słowa kluczowe:

zaćma, ból, odczuwanie bólu, lęk, fakoemulsyfikacja, operacja zaćmy.