

Zez zbieżny do dali. Diagnostyka różnicowa i metody postępowania

Distance Esotropia. Differential Diagnosis and Methods of Management

Aleksandra Budelewska¹, Piotr Loba^{1,2}

¹ Centrum Mikrochirurgii Oka Laser w Warszawie

² Zakład Patofizjologii Widzenia Obuoczniego i Leczenia Zeza I Katedry Chorób Oczu Uniwersytetu Medycznego w Łodzi
Kierownik: dr hab. n. med., prof. UMED Piotr Loba

Summary:

Primary divergence insufficiency is a generally benign condition affecting the elderly. Many patients have spontaneous resolution of the double vision within few months, so it is advisable to postpone strabismus surgery and follow the patient. It is also important to perform a differential diagnosis in order to rule out other causes of distance esotropia. For this purpose, a thorough history and careful ophthalmologic/strabologic examination should be performed. Clinical neurological evaluation is an effective tool to distinguish between patients with primary and secondary forms of divergence insufficiency. When making the diagnosis, special attention should be paid to the presence of nystagmus and the role of neuroimaging studies.

Key words:

strabology, distance esotropia, primary divergence insufficiency, secondary divergence insufficiency, diplopia.

Słowa kluczowe:

strabologia, zez zbieżny do dali, pierwotna niedomoga dywergencji, wtórna niedomoga dywergencji, dwojenie.